

## Durham Research Online

---

### Deposited in DRO:

08 May 2015

### Version of attached file:

Accepted Version

### Peer-review status of attached file:

Peer-reviewed

### Citation for published item:

Yang, S. and Fitches, E. and Pyati, P. and Gatehouse, J.A. (2015) 'Effect of insecticidal fusion proteins containing spider toxins targeting sodium and calcium ion channels on pyrethroid-resistant strains of peach-potato aphid (*Myzus persicae*).', *Pest management science.*, 71 (7). pp. 951-956.

### Further information on publisher's website:

<http://dx.doi.org/10.1002/ps.3872>

### Publisher's copyright statement:

This is the accepted version of the following article: Yang, S., Fitches, E., Pyati, P. and Gatehouse, J. A. (2014), Effect of insecticidal fusion proteins containing spider toxins targeting sodium and calcium ion channels on pyrethroid-resistant strains of peach-potato aphid (*Myzus persicae*). *Pest Management Science*, 71(7): 951-956, which has been published in final form at <http://dx.doi.org/10.1002/ps.3872>. This article may be used for non-commercial purposes in accordance With Wiley Terms and Conditions for self-archiving.

### Additional information:

## Use policy

---

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Effect of insecticidal fusion proteins containing spider toxins targeting sodium and calcium ion channels on pyrethroid-resistant strains of peach-potato aphid (*Myzus persicae*)

**Sheng Yang<sup>1</sup>, Elaine Fitches<sup>2</sup>, Prashant Pyati<sup>1</sup> and John A. Gatehouse<sup>1\*</sup>**

1 School of Biological and Biomedical Sciences, Durham University, South Road  
Durham DH1 3LE, UK

2 Fera (Food and Environment Research Agency), Sand Hutton, York, YO41 1LZ, UK

\* Corresponding author

Phone; +44 191 334 1264

FAX; +44 191 334 1201

E-mail; J.A.Gatehouse@durham.ac.uk

**Abstract:**

**BACKGROUND:** The recombinant fusion proteins PI1a/GNA and Hv1a/GNA contain the spider venom peptides  $\delta$ -amaurobitoxin-PI1a or  $\omega$ -hexatoxin-Hv1a respectively, linked to snowdrop lectin (GNA). PI1a targets receptor site 4 of insect voltage-gated sodium channels (NaCh) while Hv1a targets voltage-gated calcium channels. Insecticide-resistant strains of peach-potato aphid (*Myzus persicae*) contain mutations in NaCh. The pyrethroid-resistant "*kdr*" (794J) and "*super-kdr*" (UKO) strains contain mutations at residues L1014 and M918 in the channel  $\alpha$ -subunit respectively, while the "*kdr+super-kdr*" strain (4824J), insensitive to pyrethroids, contains mutations at both L1014 and M918.

**RESULTS:** PI1a/GNA and Hv1a/GNA fusion proteins have estimated  $LC_{50}$  values of 0.35 and 0.19 mg ml<sup>-1</sup> when fed to wild-type *M. persicae*. For insecticide-resistant aphids,  $LC_{50}$  for the PI1a/GNA fusion protein increased by 2- to 6-fold, correlating with pyrethroid resistance (wild-type < *kdr* < *super-kdr* < *kdr+super-kdr* strains). In contrast,  $LC_{50}$  for the Hv1a/GNA fusion protein showed limited correlation with pyrethroid resistance.

**CONCLUSION:** Mutations in the sodium channel in pyrethroid-resistant aphids also protect against a fusion protein containing a sodium channel-specific toxin, despite differences in ligand-channel interactions, but do not confer resistance to a fusion protein targeting calcium channels. The use of fusion proteins with differing targets could play a role in managing pesticide resistance.

**Key words:** biopesticide; insecticide resistance; Homoptera / Hemiptera; voltage-gated ion channels; fitness cost

## 1 INTRODUCTION

The peach-potato aphid, *Myzus persicae* (Sulzer) (Hemiptera: Aphididae), is a serious worldwide insect pest of agricultural and horticultural crops, which, through its sap-sucking feeding habit, can transmit viral diseases.<sup>1</sup> Pyrethroids are a major class of insecticides used to control this pest, but populations of *M. persicae* can rapidly develop resistance to pyrethroids, leading to increased economic loss to agricultural producers.<sup>2</sup> Pyrethroids target the insect voltage-gated sodium channel, a large transmembrane protein composed of a single 260kDa polypeptide (the alpha subunit), which contains four repeating and homologous domains (I–IV), with each domain being constituted by six hydrophobic transmembrane segments (S1–S6).<sup>3</sup> The insect sodium channel is similar in structure to the vertebrate sodium channel, containing different allosterically coupled receptor-binding sites for various neurotoxicants, but the two types of channel are distinguishable in the pharmacology. Therefore insecticides such as pyrethroids can be specific for insect sodium channels, showing no effect on mammals.<sup>4,5</sup>

Pyrethroids are hydrophobic compounds, and are thought to bind to the lipid-exposed interface formed by helices IIS6, IIS5, linker helix IIS4-IIS5 and the IS4-IS5 linker,<sup>6,7</sup> affecting the functional properties of the sodium channel. By preventing closure of the sodium channel, pyrethroids cause paralysis in insects.<sup>5</sup> However, with the extensive use of pyrethroids, many insects have developed resistance to these insecticides, associated with mutations in the sodium channel. The pyrethroid resistance shown by *M. persicae* is typical of that seen in many species.<sup>8-10</sup> In aphids carrying the *kdr* mutation, there is a leucine to phenylalanine substitution (L1014F) within segment 6 of domain II (IIS6) of the channel protein,<sup>11</sup> which confers an intermediate level of resistance to pyrethroids. In aphids carrying the *super-kdr* site mutation, there is an additional methionine- to-threonine substitution (M918T) in the linker between segment 4 and segment 5 of domain II (IIS4-IIS5 linker) of the sodium channel protein,<sup>8</sup> which makes *M. persicae* highly resistant to pyrethroids. Data

presented by Eleftherianos et al.<sup>1</sup> shows that whereas the EC<sub>50</sub> for a typical pyrethroid insecticide on wild-type *M. persicae* is in the range 0.5 - 2.8 ppm, a homozygous *kdr* mutation increases the EC<sub>50</sub> by 20-75 fold, and a heterozygous *kdr*+*super-kdr* mutation increases resistance by 100-500 fold. The emergence of insecticide resistance is one factor driving a need for new specific environmentally benign pesticides, which could be used in strategies to manage resistance to chemicals like pyrethroids more effectively.

Spider toxin peptides have been suggested as environmentally friendly biopesticides. Toxins have been isolated from a range of arachnids, and most are small cysteine-rich proteins that principally target neuronal ion channels to cause paralysis of the spider's prey.<sup>4,12</sup> Toxins can be selected that are insect-specific, and have no effects on members of other taxons. This advantage would make them ideal candidates for use in pest control and crop protection, if a suitable delivery system which would get around the problem of toxicity being dependent on injection into the body fluid of the pest could be devised.<sup>13</sup> Recombinant fusion proteins, containing insecticidal peptides or proteins fused to a "carrier" protein are a method, which gives oral toxicity to neuroactive toxins.<sup>14,15</sup> The carrier protein transports the insecticidal peptide or protein across the insect gut epithelium into the haemolymph, from which it can access the central nervous system (CNS), which is the site of action. The mannose-specific lectin from snowdrop (*Galanthus nivalis* agglutinin: GNA), which has been shown to transport peptides into the insect haemolymph, is currently being used for making fusion proteins. Fusion proteins containing GNA as a carrier possess good stability towards proteolysis in the insect gut and high toxicity.<sup>16</sup>

δ-Amaurobitoxins, or δ-palutoxins, from the spider *Pireneitega luctuosus*, are a family of four similar 36-37 residue peptides containing 8 cysteine residues which are disulphide-linked to form a cysteine knot motif. PI1a is specific for insect sodium channels, causing paralysis, and has no adverse effects when injected into mice.<sup>17</sup> The toxin acts by binding to receptor site 4 in the sodium channel protein, which involves

the extracellular loops of S1-S2, S3-S4 of domain II.<sup>18</sup> It affects the functional properties of the sodium channel  $\alpha$  subunit by shifting the voltage dependence of activation, resulting in paralysis; the effect is similar to that produced by pyrethroids.<sup>5</sup> A PI1a/GNA fusion protein has been shown to be an effective oral insecticide towards insects of different orders, including aphids.<sup>19</sup>

Hv1a is a family member of insecticidal neurotoxins, which possess 36–37 residues, from the Australian funnel web spider *Hadronyche versuta*.<sup>20</sup> Hv1a arrests insect voltage-gated calcium channels and has no negative effects on mammals.<sup>21-23</sup> Hv1a contains three disulfide bonds which shape an inhibitor cystine knot motif, which confers chemical and thermal stability and resistance to proteases.<sup>24,25</sup> The highly conserved C-terminal  $\beta$  hairpin of Hv1a contains the key residues for insecticidal activity.<sup>20</sup> An Hv1a/GNA fusion protein has been described previously, and its oral toxicity towards insects has been demonstrated.<sup>16</sup>

The present paper compares the toxicity of PI1a/GNA and Hv1a/GNA fusion proteins towards wild-type and pyrethroid-resistant strains of *M. persicae*, and shows that although the toxicity of PI1a/GNA is reduced by the *kdr* and *super-kdr* mutations in the sodium channel, it retains some activity. However, the mutations confer no resistance to Hv1a/GNA targeting calcium channels. This residual high insecticidal activity makes Hv1a/GNA a potential biopesticide for controlling pyrethroid-resistant aphids.

## 2 MATERIALS AND METHODS

### 2.1 Materials

Chemicals and reagents were of analytical grade and were supplied by Sigma or BDH Chemical Company otherwise unless stated. Restriction enzymes and other molecular biology reagents were supplied by Fermentas. A double stranded DNA incorporating a sequence encoding the mature PI1a toxin (P83256), with codons optimised for

expression in *Pichia pastoris*, was designed by the authors, synthesized and supplied by ShineGene Molecular Biotech, Inc. (Shanghai 201109, China; <http://www.synthesiscgene.com/>). Other oligonucleotides required for cloning were supplied by Sigma Chemical Co. Recombinant snowdrop lectin was produced by the authors by expression in *Pichia pastoris*, as described by Baumgartner et al. (2004).<sup>26</sup>

The mutant strains of peach-potato aphid, *Myzus persicae* (Sulzer) (Hemiptera: Aphididae) were kindly provided by Prof. Linda M. Field (Department of Biological Chemistry and Crop Protection, Rothamsted Research, UK). Strain 4106A has no mutation ("wild type"). Strain 794J is homozygous for the mutation L1014F (*kdr*), and is resistant to pyrethroids. Strain UKO is homozygous for the mutation M918L (*super-kdr*), and shows enhanced resistance to pyrethroids. 4824J is homozygous for L1014F (*kdr*) and M918T (*super-kdr*), and shows immunity to pyrethroids.<sup>1</sup> Aphids were cultured on fresh Chinese Leaf under conditions of 12h light, 12h dark, 18°C, 70% relative humidity.

## **2.2 Production of PI1a/GNA and Hv1a/GNA fusion proteins**

Assembly of expression constructs encoding PI1a, PI1a/GNA and GNA and expression of the recombinant proteins in the yeast *Pichia pastoris* have been described elsewhere.<sup>19</sup> The fusion proteins, which contained C-terminal (His)<sub>6</sub> tags, were purified by metal affinity chromatography, dialysed and lyophilised as previously described.<sup>14-16</sup> Expression constructs for Hv1a and Hv1a/GNA and production of recombinant proteins have also been described previously;<sup>16</sup> the constructs used to express Hv1a and Hv1a/GNA for this paper were modified by inclusion of a predicted pro-region for the toxin.<sup>27</sup> Other recombinant proteins were produced as previously described.<sup>15</sup> Purified proteins were analysed by SDS-PAGE for quantitation by comparison to standards run on the same gel; proteins were also quantitated by using the BCA assay, and by absorbance.

## **2.3 Bioassays on peach-potato aphid**

Bioassay of aphids using liquid artificial diet was carried out as described by Prosser and Douglas<sup>28</sup>. Adult aphids were transferred to control liquid diet, acclimatised for 24h, and then neonate nymphs produced over the following 24h were transferred to experimental diets, and allowed to develop to adult stage (8-9 days). 20 individuals per treatment were used to perform the bioassays. Each assay was repeated 3 times. Mortality was observed daily, and assays were continued until control aphids started to produce nymphs. Nymphs were not counted but the presence or absence of progeny was recorded. Effects of treatments on aphid growth were assessed by using Image J Software to measure insect length.

## **2.4 Statistical analysis**

Mortality data were analysed using survival curves, with a Kaplan-Meier test to evaluate significance of differences (Origin 8.5 software). ANOVA analysis (with Bonferroni-Dunn post-hoc tests) was carried out to determine any significant differences between treatments in size parameters measured. Differences between treatments were considered significant at a probability level  $p < 0.05$ .  $LC_{50}$  values for different treatments were estimated by taking survival data for diets containing different concentrations of fusion proteins (over a range of 0.125 - 2.0 mg ml<sup>-1</sup>) and fitting data points to a sigmoidal dose-response curve by non-linear regression (Prism v. 5 software).

## **3 RESULTS**

### **3.1 Toxicity of separate components of fusion proteins**

Effects of toxins and GNA components of insecticidal fusion proteins on the strains of peach-potato aphids (794J, UKO, 4824J and 4106A) were determined by bioassays in which components were fed separately in liquid diet from neonate nymphs. Concentrations were chosen to be equivalent to 1 mg/ml fusion protein. Results are



shown in Fig. 1. None of the treatments caused more than 30% mortality over a 7-day period of development against a background of no mortality in aphids on control diet; survival analysis showed that most differences to control were not significant (effect on survival by difference in survival curve;  $p > 0.05$ ). The GNA carrier protein showed significant effects on *M. persicae* survival (difference in survival curve;  $p < 0.05$ ), in agreement with previous reports that this protein is weakly insecticidal towards aphids<sup>29</sup>; it also caused growth retardation at the beginning in the bioassays, although aphids were able to recover from the effects and produced nymphs. There were no significant differences in the effects of GNA between aphid strains. At the concentrations used, the Hv1a toxin showed significant effects on *M. persicae* (30% mortality after 7 days; effect on survival by difference in survival curve  $p < 0.05$ ), whereas PI1a did not have a significant effect, although both toxins have been shown previously to have some effect on aphids when fed in diet. Once again, no significant differences between aphid strains were observed in these assays. These data confirm previous observations that the separate components of insecticidal fusion proteins have only limited insecticidal effects when fed to *M. persicae*.

### 3.2 Toxicity of PI1a/GNA recombinant fusion protein

Purified recombinant PI1a/GNA fusion protein was fed to each *M. persicae* strain at a range of concentrations, and survival curves were plotted for all treatments. Results for feeding at 1 mg ml<sup>-1</sup> are shown in Fig 2A. At this level, the fusion protein caused complete mortality to strain 4106A (*wild-type*) after 7 days, but not in any of the insecticide resistant strains, even after 11 days. The survival curves show significant differences between strains 4106A (*wild-type*), 794J (*kdr*) and UKO (*super-kdr*) and the controls not fed fusion protein ( $\geq 90\%$  survival) ( $p < 0.05$ ), confirming the insecticidal activity of the treatment. However, the survival curve for strain 4824J (*kdr + super-kdr*: 90% survival over the assay) fed PI1a/GNA at 1 mg ml<sup>-1</sup> is not significantly different to that for aphids fed control diet containing no fusion protein ( $p < 0.05$ ). Survival curves

for strains 794J (*kdr*) and UKO (*super-kdr*), which both show 40% survival over the assay, differ significantly from controls, from wild-type survival, and from strain 4824J survival ( $p < 0.05$ ). Growth retardation was observed in all aphids exposed to fusion proteins, but was least in strain 4824J (Fig. 2B), where aphids were able to produce nymphs during the assay period, as did the controls. No other aphid strain exposed to treatment was able to produce nymphs. The data demonstrate a differential effect of the fusion protein on the different aphid strains, with *wild-type* strains fully susceptible to the toxin at this concentration, whereas the *kdr* and *super-kdr* strains are partially tolerant, and the *kdr + super-kdr* strain is almost completely tolerant.

By analysing survival curves for aphids exposed to different concentrations of PI1a/GNA,  $LC_{50}$  values for the different strains could be deduced. The values obtained range from 0.35 to 1.76 mg ml<sup>-1</sup>, and are shown in Table 1. There is a strong correlation between insecticide resistance of aphid strains and the estimated  $LC_{50}$  values; wild-type susceptible aphids have the lowest  $LC_{50}$ , and the order of insecticide tolerance (*wild-type* < *kdr* < *super-kdr* < *kdr + super-kdr*) is reflected in the  $LC_{50}$  values (*wild-type* < *kdr* < *super-kdr* < *kdr + super-kdr*). The *kdr + super-kdr* strain 4824J has an estimated  $LC_{50}$  of 1.76 mg ml<sup>-1</sup> for PI1a/GNA; recombinant protein at 2.0 mg ml<sup>-1</sup> caused significant effects on survival, and treatment with 2.5 or 3.0 mg ml<sup>-1</sup> of PI1a/GNA resulted in complete mortality (Fig. 2C).

### 3.3 Toxicity of Hv1a/GNA recombinant fusion protein

An insecticidal fusion protein containing the calcium-channel specific toxin Hv1a was used as a control to identify non-specific effects on sensitivity towards insecticidal compounds in the pyrethroid-resistant *M. persicae* strains. Purified recombinant Hv1a/GNA fusion protein was fed to each strain at a range of concentrations, and survival curves were plotted for all treatments. Results for feeding at 1 mg ml<sup>-1</sup> are shown in Fig.3A. Hv1a/GNA fusion protein at this concentration caused complete mortality to strains 4106A (*wild-type*) and UKO (*super-kdr*) after 6 days, and to strains

794J (*kdr*) and 4824J (*kdr* + *super-kdr*) after 9 days. The survival curves show significant differences between all strains fed fusion protein and the controls not fed fusion protein (100% survival over 11 days) ( $p < 0.05$ ), in agreement with previous assays showing that this fusion protein is insecticidal. Growth retardation was observed in all aphids exposed to fusion proteins (Fig 3B), and no aphids exposed to treatment were able to produce nymphs. Comparison of individual survival curves when Hv1a/GNA was fed at  $1 \text{ mg ml}^{-1}$  suggested that strain 4824J (*kdr* + *super-kdr*) was more tolerant to Hv1a/GNA than *wild-type* aphids (strain 4106A), (difference between survival curves at  $p < 0.05$ ) but that other differences were not significant. Assays at other concentrations of Hv1a/GNA did not give consistently significant differences between treatments, although the *wild-type* strain always showed greater susceptibility to the fusion protein than the pyrethroid-resistant strains.

LC<sub>50</sub> values for Hv1a/GNA in the different aphid strains were deduced by analysis of survival curves for aphids exposed to different concentrations fusion protein. The values obtained range from 0.19 to 0.28  $\text{mg ml}^{-1}$ , and are shown in Table 1. The estimated LC<sub>50</sub> values show no significant differences between any of the aphid strains although the *wild-type* strain, 4106A, has a lowest LC<sub>50</sub> value. The uncertainties in estimated LC<sub>50</sub> values are relatively large compared to the differences, but the fitted dose-response curve for the *wild-type* strain differs significantly from the other curves ( $p < 0.05$ ), supporting the conclusion that this strain is more susceptible to Hv1a/GNA.

#### 4 DISCUSSION

The insect sodium channel is a major target for conventional pesticides, such as pyrethroids. The PI1a toxin, which acts on the same target, could represent a novel type of insecticidal component as a substitute to pyrethroids. The mode of binding of this toxin would be expected to differ significantly from binding a small molecule channel blocker like a pyrethroid, with contacts between the toxin and the channel potentially extending over a wider area. However, PI1a/GNA fusion protein exhibits

reduced toxicity towards pyrethroid-resistant peach-potato aphid (*Myzus persicae*) strains, showing that the mutations, which remove sensitivity to pyrethroids, also affect the binding of PI1a. The mutations which give pyrethroid sensitivity are in domain II of the sodium channel, with the mutation at L1014 in helix S6 and the mutation at M918 in the linker between helices S4-S5. Changes to the spatial structure of domain II as a result of these mutations presumably also disturb the binding of PI1a to receptor site 4, in domain II. However, although the bioassays show that mutations in domain II of the insect sodium channel affect the insecticidal activity of the PI1a/GNA fusion protein, some toxicity is still observed, with a higher concentration of fusion protein required to cause mortality in the pyrethroid-resistant *kdr* and *super-kdr* strains. This result implies that either some interactions still exist between PI1a and domain IIS6 or domain IIS4-S5 linker of the mutated sodium channel, or that PI1a also binds to other sites on the sodium channel to cause inactivation. The extracellular loops of IIS1-S2, IIS3-S4 are thought to be the main binding sites of PI1a, which are distinct from the pyrethroid binding site but contribute to receptor site 4 for toxins. The change in the spatial structure of domain II as a result of the *kdr* and *super-kdr* mutations may have a relatively small effect on toxin binding in the interaction between PI1a and the sodium channel but may prevent the toxin inactivating the channel. The greater effect on channel structure caused by combining the mutations at L1014 and M918 would be expected to affect PI1a binding more than single mutations, in agreement with the lack of sensitivity to PI1a/GNA shown by aphid strain 4824J.

As expected, when fusion protein containing the calcium channel-specific toxin Hv1a is fed to aphids, there is no evidence for significant differential sensitivity between insecticide-resistant aphid strains, since the strains differ in mutations to the sodium channel. However, the observation that wild-type aphids are more susceptible to this toxin is unexpected. Mutations in sodium channels present in strains 794J, UKO and 4824J would be expected to result in a fitness cost to *M. persicae*, similar to that observed both for other insect-resistant aphids of this species,<sup>30</sup> and for other insect

species (e.g. when comparing insecticide-resistant and insecticide-susceptible German cockroaches, *Blattella germanica*<sup>31</sup>) A fitness cost for insecticide resistance can be inferred in *M. persicae* from population data; if there were no fitness cost, the population of resistant *M. persicae* should be much larger than wild type before selection occurs.<sup>32</sup> The fitness cost would be expected to make insecticide-resistant strains of *M. persicae* more susceptible to Hv1a/GNA, but this is not the case. Possibly, other changes to the phenotype of insecticide-resistant aphids are affecting susceptibility to this fusion protein; a transcriptomic study<sup>33</sup> has suggested that insecticide resistance in *M. persicae* is complex, and involves a broad array of resistance mechanisms. The present results support that conclusion.

The *kdr* strain of *M. persicae* is resistant to all pyrethroids, showing 23-to 73-fold increased resistance<sup>1</sup> and the *kdr* +*super-kdr* strain is virtually immune to all the pyrethroids.<sup>34</sup> A fusion protein containing the sodium-channel specific PI1a toxin can cause 100% mortality towards pyrethroid-resistant aphids containing a single mutation in the sodium channel if administered at concentrations increased only 3-fold, but is not effective towards aphids containing a double mutation in the sodium channel. However, insecticide-resistant aphids are still sensitive towards a calcium channel-specific toxin, albeit at higher doses than wild-type aphids. These experiments demonstrate the potential for fusion protein-based biopesticides to complement existing pesticides, and to be used in the management of insecticide-resistant insect strains; the Hv1a/GNA fusion protein is currently undergoing trials leading to commercial use as a biopesticide.

## Acknowledgements

Work described in this paper has been carried out with funding from the Technology Strategy Board (TSB; TS/I000690/1) and BBSRC China Partnership Scheme (BB/G530409), which is acknowledged with gratitude. SY received support for his Ph.D. programme jointly from the China Scholarship Scheme, and a Durham University Doctoral Studentship; this funding is also gratefully acknowledged.

## References

1. Eleftherianos I, Foster SP, Williamson MS and Denholm I, Characterization of the M918T sodium channel gene mutation associated with strong resistance to pyrethroid insecticides in the peach-potato aphid, *Myzus persicae* (Sulzer). *Bull Entomol Res* **98**: 183-191 (2008).
2. McCaffery A and Nauen R, The insecticide resistance action committee (IRAC): public responsibility and enlightened industrial self-interest. *Outlooks on Pest Management* **17**: 11-14 (2006).
3. Catterall WA, From ionic currents to molecular mechanisms: the structure and function of voltage-gated sodium channels. *Neuron* **26**: 13-25 (2000).
4. Vassilevski AA, Kozlov SA and Grishin EV, Molecular diversity of spider venom. *Biochemistry (Mosc)* **74**: 1505-1534 (2009).
5. Zlotkin E, The insect voltage-gated sodium channel as target of insecticides. *Annu Rev Entomol* **44**: 429-45 (1999).
6. Du Y, Lee J, Nomura Y, Zhang T, Zhorov B and Dong K, Identification of a cluster of residues in transmembrane segment 6 of domain III of the cockroach sodium channel essential for the action of pyrethroid insecticides. *Biochem J* **419**:377-385 (2009).
7. Du Y, Nomura Y, Satar G, Hu Z, Nauen R, He SY, Zhorov BS and Dong K, Molecular evidence for dual pyrethroid-receptor sites on a mosquito sodium channel. *Proc Natl Acad Sci U S A* **110**: 11785-11790 (2013).
8. Williamson MS, Martinez-Torres D, Hick CA and Devonshire AL, Identification of mutations in the housefly *para*-type sodium channel gene associated with knockdown resistance (*kdr*) to pyrethroid insecticides. *Mol Gen Genet* **252**: 51-60 (1996).
9. Martinez-Torres D, Chandre F, Williamson MS, Darriet F, Bergé JB, Devonshire AL, Guillet P, Pasteur N and Pauron D, Molecular characterization of pyrethroid

308 knockdown resistance (*kdr*) in the major malaria vector *Anopheles gambiae* ss.  
309 *Insect Mol Biol* **7**: 179-184 (1998).

310 10. Kranthi KR, Jadhav DR, Kranthi S, Wanjari RR, Ali SS and Russell DA, Insecticide  
311 resistance in five major insect pests of cotton in India. *Crop Prot* **21**: 449-460  
312 (2002).

313 11. Miyazaki M, Ohyama K, Dunlap DY and Matsumura F, Cloning and sequencing of  
314 the *para*-type sodium channel gene from susceptible and *kdr*-resistant German  
315 cockroaches (*Blattella germanica*) and house fly (*Musca domestica*). *Mol Gen*  
316 *Genet* **252(1-2)**: 61-68 (1996).

317 12. Rash LD and Hodgson WC, Review: pharmacology and biochemistry of spider  
318 venoms. *Toxicon* **40**: 225–254 (2002).

319 13. Whetstone PA and Hammock BD, Delivery methods for peptide and protein  
320 toxins in insect control. *Toxicon* **49**: 576-596 (2007).

321 14. Fitches E, Audsley N, Gatehouse JA and Edwards JP, Fusion proteins  
322 containing neuropeptides as novel insect control agents: snowdrop lectin delivers  
323 fused allatostatin to insect haemolymph following oral ingestion. *Insect Biochem*  
324 *Mol Biol* **32**: 1653-1661 (2002).

325 15. Fitches E, Edwards MG, Mee C, Grishin E, Gatehouse AM, Edwards JP and  
326 Gatehouse JA, Fusion proteins containing insect-specific toxins as pest control  
327 agents: snowdrop lectin delivers fused insecticidal spider venom toxin to insect  
328 haemolymph following oral ingestion. *J Insect Physiol* **50**: 61-71 (2004).

329 16. Fitches EC, Pyati P, King GF and Gatehouse JA, Fusion to snowdrop lectin  
330 magnifies the oral activity of insecticidal  $\omega$ -hexatoxin-Hv1a peptide by enabling its  
331 delivery to the central nervous system. *PLoS One* **7**: e39389 (2012).

332 17. Corzo G., Escoubas P, Stankiewicz M, Pelhate M, Kristensen CP and Nakajima T,  
333 Isolation, synthesis and pharmacological characterization of  
334  $\delta$ -palutoxins IT, novel insecticidal toxins from the spider *Paracoelotes luctuosus*  
335 (*Amaurobiidae*). *Eur J Biochem* **267**: 5783-5795 (2000).

- 336 18. Cestèle S and Catterall WA, Molecular mechanisms of neurotoxin action on  
337 voltage-gated sodium channels. *Biochimie* **82**: 883-892 (2000).
- 338 19. Yang S, Pyati P, Fitches E and Gatehouse JA, A recombinant fusion protein  
339 containing a spider toxin specific for the insect voltage-gated sodium ion channel  
340 shows oral toxicity towards insects of different orders. *Insect Biochem Mol* **47**: 1-  
341 11 (2014).
- 342 20. Tedford HW, Fletcher JI and King GF, Functional significance of the  $\beta$ -hairpin in  
343 the insecticidal neurotoxin  $\omega$ -atracotoxin-Hv1a. *J Biol Chem* **276**: 26568-26576  
344 (2001).
- 345 21. Fletcher JI, Smith R, O'Donoghue SI, Nilges M, Connor M, Howden ME, Christie  
346 MJ and King GF, The structure of a novel insecticidal neurotoxin,  $\omega$ -atracotoxin-  
347 HV1, from the venom of an Australian funnel web spider. *Nat Struct Biol* **4**: 559-  
348 566 (1997).
- 349 22. Tedford HW, Gilles N, Ménez A, Doering CJ, Zamponi GW and King GF, Scanning  
350 mutagenesis of  $\omega$ -atracotoxin-Hv1a reveals a spatially restricted epitope that  
351 confers selective activity against insect calcium channels. *J Biol Chem* **279**:  
352 44133-44140 (2004).
- 353 23. Chong Y, Hayes JL, Sollod B, Wen S, Wilson DT, Hains PG, Hodgson WC, Broady  
354 KW, King GF and Nicholson GM, The  $\omega$ -atracotoxins: selective blockers of insect  
355 M-LVA and HVA calcium channels. *Biochem Pharmacol* **74**: 623-638 (2007).
- 356 24. King GF, Tedford HW and Maggio F, Structure and function of insecticidal  
357 neurotoxins from Australian funnel-web spiders. *Toxin Rev* **21(4)**: 361-389 (2002).
- 358 25. Saez NJ, Senff S, Jensen JE, Er SY, Herzig V, Rash LD and King GF, Spider-  
359 venom peptides as therapeutics. *Toxins* **2**: 2851-2871 (2010).
- 360 26. Baumgartner P, Harper K, Raemaekers RJM, Durieux A, Gatehouse AMR, Davies  
361 HV and Taylor MA, Large-scale production and purification of recombinant  
362 *Galanthus nivalis* agglutinin (GNA) expressed in the methylotrophic yeast *Pichia*  
363 *pastoris*. *Biotechnol Letts* **25**: 1281-1285 (2004).



27. Gatehouse JA, Yang S, Pyati P and Fitches EC, Pesticidal Fusion Protein Improvements. *UK Patent Applic. No.* 1321938.1 (2013).
28. Prosser WA and Douglas AE, A test of the hypotheses that nitrogen is upgraded and recycled in an aphid (*Acyrtosiphon pisum*) symbiosis. *J Insect Physiol* **38**: 93-99 (1992).
29. Down RE, Fitches EC, Wiles DP, Corti P, Bell HA, Gatehouse JA and Edwards JP, Insecticidal spider venom toxin fused to snowdrop lectin is toxic to the peach-potato aphid, *Myzus persicae* (Hemiptera: Aphididae) and the rice brown planthopper, *Nilaparvata lugens* (Hemiptera: Delphacidae). *Pest Manag Sci* **62**: 77-85 (2006).
30. Castañeda LE, Barrientos K, Cortes PA, Figueroa CC, Fuentes-Contreras E, Luna-Rudloff M, Silva AX and Bacigalupe LD, Evaluating reproductive fitness and metabolic costs for insecticide resistance in *Myzus persicae* from Chile. *Physiol Entomol* **36**: 253–260.
31. Ang LH and Lee CY, Absence of a fitness penalty in insecticide-resistant German cockroaches, *Blattella germanica* (L.)(Dictyoptera: Blattellidae). *International Journal of Pest Management* **57**: 195-204 (2011).
32. Fenton B, Margaritopoulos JT, Malloch GL and Foster S P, Micro-evolutionary change in relation to insecticide resistance in the peach–potato aphid, *Myzus persicae*. *Ecol Entomol* **35**: 131-146 (2010).
- 33 Silva AX, Jander G, Samaniego H, Ramsey JS and Figueroa CC, Insecticide resistance mechanisms in the green peach aphid *Myzus persicae* (Hemiptera: Aphididae) I: A Transcriptomic Survey. *PLoS ONE* **7**: e36366 (2012).
34. Anstead JA1, Williamson MS, Eleftherianos I and Denholm I, High-throughput detection of knockdown resistance in *Myzus persicae* using allelic discriminating quantitative PCR. *Insect Biochem Mol Biol* **34**: 871-877 (2004).

**Table 1.** Estimated LC<sub>50</sub> values for fusion proteins towards wild-type and pyrethroid tolerant strains of *M. persicae*. Values were calculated from dose-response curves fitted to survival data after 9 days' exposure to diets containing fusion proteins at varying concentrations.

Genotype (Strain)	LC <sub>50</sub> (mg ml <sup>-1</sup> ) PI1a/GNA	LC <sub>50</sub> (mg ml <sup>-1</sup> ) Hv1a/GNA
4106A (wild type)	0.35	0.19
794J ( <i>kdr</i> )	0.60	0.28
UKO ( <i>super-kdr</i> )	0.83	0.25
4824J ( <i>kdr</i> + <i>super-kdr</i> )	1.76	0.20

## Figure Legends

**Figure 1.** Toxicity of fusion protein components towards *M. persicae*. Graph shows survival after 7 days of pyrethroid-tolerant *M. persicae* strains (794J, *kdr*, UKO, *super-kdr*, and 4824J, *kdr+super-kdr*) and wild type 4106A strain after feeding artificial diet containing 0.4 mg ml<sup>-1</sup> PI1a, 0.46 mg ml<sup>-1</sup> Hv1a or 0.6 mg ml<sup>-1</sup> GNA. Survival on control diet was 100% for all aphid strains over this interval. n = 20 aphids per replicate.

## Figure 2.

**(A):** Toxicity of PI1a/GNA fusion protein towards *M. persicae*. Graph shows survival curves of pyrethroid-tolerant *M. persicae* strains (794J, *kdr*, UKO, *super-kdr*, and 4824J, *kdr+super-kdr*), and wild type 4106A strain fed PI1a/GNA at 1 mg ml<sup>-1</sup>. All aphid strains on control diet showed survival similar to that presented for 4106A strain. n = 20 aphids per replicate.

**(B):** Growth suppression by PI1a/GNA fusion protein. Graph shows lengths of aphid strains 794J, *kdr*, UKO, *super-kdr*, and 4824J, *kdr+super-kdr* and 4106A (wild type) from neonate to adult (9 days) after feeding on artificial diet containing 1mg/ml PI1a/GNA (n=3 per treatment). 100 % mortality for strain 4106A prevented analysis for day 9. Data for strain 4842J fed on control diet is shown, but all aphid strains fed on control diet were of comparable size at each time point.

**(C):** Dose-response effects of PI1a/GNA. Graph shows survival curves of 4824J (*kdr+super-kdr*) *M. persicae* strain fed diets containing different concentrations of PI1a/GNA in the range 0 - 3.0 mg ml<sup>-1</sup>. n=20 aphids per replicate.

## Figure 3.

**(A):** Toxicity of Hv1a/GNA fusion protein towards *M. persicae*. Graph shows survival of pyrethroid-tolerant *M. persicae* strains (794J, *kdr*, UKO, *super-kdr*, and 4824J, *kdr+super-kdr*) and wild type 4106A strain fed on diet containing 1 mg ml<sup>-1</sup> Hv1a/GNA. All aphid strains on control diet showed survival similar to that presented for 4106A

428 strain. n = 20 aphids per replicate.

429 **(B):** Growth suppression by PHv1a/GNA fusion protein. Graph shows lengths of aphid  
430 strains 794J, *kdr*, UKO, *super-kdr*, and 4824J, *kdr+super-kdr* and 4106A (wild type)  
431 from neonate to adult after feeding on artificial diet containing 1mg/ml Hv1a/GNA (n=3  
432 per treatment). 100 % mortality for strains UKO, 4824J and 4106A prevented analysis  
433 for day 9. Data for strain 4842J fed on control diet is shown, but all aphid strains fed on  
434 control diet were of comparable size at each time point.

435

436

437

438